



U production at NLO and beyond

J.P. Lansberg

► To cite this version:

J.P. Lansberg. U production at NLO and beyond. Workshop on Charmonium production and decays : new results and perspectives, Mar 2013, Orsay, France. in2p3-00856690

HAL Id: in2p3-00856690

<https://hal.in2p3.fr/in2p3-00856690>

Submitted on 2 Sep 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Υ production at NLO and beyond

J.P. Lansberg

IPN Orsay – Paris-Sud U. –CNRS/IN2P3

**Workshop on Charmonium production and decays :
new results and perspectives**

LAL Orsay

March 6-8, 2013



Part I

(Single) Υ production in pp

Reminder: QCD corrections for Υ at the Tevatron



Reminder: QCD corrections for Υ at the Tevatron

J.Campbell, F. Maltoni, F. Tramontano, Phys.Rev.Lett. 98:252002,2007
 P.Artoisenet, J.Campbell, JPL, F.Maltoni, F. Tramontano, Phys. Rev. Lett. 101, 152001 (2008)
 CDF PRL 88 (2002) 161802; LHCb EPJC 72 (2012) 2025



Reminder: QCD corrections for Υ at the Tevatron

J.Campbell, F. Maltoni, F. Tramontano, Phys.Rev.Lett. 98:252002,2007
 P.Artoisenet, J.Campbell, JPL, F.Maltoni, F. Tramontano, Phys. Rev. Lett. 101, 152001 (2008)
 CDF PRL 88 (2002) 161802; LHCb EPJC 72 (2012) 2025




Reminder: QCD corrections for Υ at the Tevatron


J.Campbell, F. Maltoni, F. Tramontano, Phys.Rev.Lett. 98:252002,2007

P.Artoisenet, J.Campbell, JPL, F.Maltoni, F. Tramontano, Phys. Rev. Lett. 101, 152001 (2008)

CDF PRL 88 (2002) 161802; LHCb EPJC 72 (2012) 2025





+ double t -channel gluon exchange at α_S^5



QCD corrections for Υ at the Tevatron & the LHC



J.Campbell, F. Maltoni, F. Tramontano, Phys.Rev.Lett. 98:252002,2007
 P.Artoisenet, J.Campbell, JPL, F.Maltoni, F. Tramontano, Phys. Rev. Lett. 101, 152001 (2008)
 CDF PRL 88 (2002) 161802; LHCb EPJC 72 (2012) 2025



$\Upsilon(3S)$: 100 % direct; $\Upsilon(2S)$: 60-70 % direct; $\Upsilon(1S)$: 50 % direct

QCD corrections for Υ at the Tevatron & the LHC



J.Campbell, F. Maltoni, F. Tramontano, Phys.Rev.Lett. 98:252002,2007
 P.Artoisenet, J.Campbell, JPL, F.Maltoni, F. Tramontano, Phys. Rev. Lett. 101, 152001 (2008)
 CDF PRL 88 (2002) 161802; LHCb EPJC 72 (2012) 2025



$\Upsilon(3S)$: 100 % direct; $\Upsilon(2S)$: 60-70 % direct; $\Upsilon(1S)$: 50 % direct



QCD corrections for Υ at the Tevatron & the LHC

J.Campbell, F. Maltoni, F. Tramontano, Phys.Rev.Lett. 98:252002,2007
 P.Artoisenet, J.Campbell, JPL, F.Maltoni, F. Tramontano, Phys. Rev. Lett. 101, 152001 (2008)
 CDF PRL 88 (2002) 161802; LHCb EPJC 72 (2012) 2025



$\Upsilon(3S)$: 100 % direct; $\Upsilon(2S)$: 60-70 % direct; $\Upsilon(1S)$: 50 % direct

Analogy with the P_T spectrum for the Z^0 boson



CSM predictions account for the P_T -integrated yield

S. J. Brodsky and JPL, PRD 81 051502 (R), 2010; JPL, PoS(ICHEP 2010), 206 (2010); NPA (2012), 10.1016/j.nuclphysa.2012.12.051

→ The **yield vs. \sqrt{s}** (here only LO curves¹)

¹NLO not stable at large \sqrt{s} (small x) and small P_T

CSM predictions account for the P_T -integrated yield

S. J. Brodsky and JPL, PRD 81 051502 (R), 2010; JPL, PoS(ICHEP 2010), 206 (2010); NPA (2012), 10.1016/j.nuclphysa.2012.12.051

→ The **yield vs. \sqrt{s}** (here only LO curves¹)

- Unfortunately, very large th. uncertainties: masses, scales (μ_R , μ_F), gluon PDFs at low x and Q^2 , ...
- Good agreement with RHIC, Tevatron and LHC data
(multiplied by a constant F^{direct})



¹NLO not stable at large \sqrt{s} (small x) and small P_T

CSM predictions account for the P_T -integrated yield

S. J. Brodsky and JPL, PRD 81 051502 (R), 2010; JPL, PoS(ICHEP 2010), 206 (2010); NPA (2012), 10.1016/j.nuclphysa.2012.12.051

→ The **yield vs. \sqrt{s}** (here only LO curves¹)

- Unfortunately, very large th. uncertainties: masses, scales (μ_R , μ_F), gluon PDFs at low x and Q^2 , ...
- Good agreement with RHIC, Tevatron and LHC data



STAR PRD 82 (2010) 012004 ; CDF PRL 88 (2002) 161802; CMS PRD 83 (2011) 112004; LHCb EPJC 72 (2012) 2025

¹NLO not stable at large \sqrt{s} (small x) and small P_T

Cross section ratio I

- Despite th. uncertainties, CSM predictions are **parameter free** !

Cross section ratio I

- Despite th. uncertainties, CSM predictions are **parameter free** !
- At LO in v^2 , one *de facto* **predicts direct** cross-section **ratios**

Cross section ratio I

- Despite th. uncertainties, CSM predictions are **parameter free** !
- At LO in v^2 , one *de facto* **predicts direct** cross-section **ratios**
- Simple ratios of **Schrödinger wave function** at the origin:

$$\frac{\sigma(\text{direct } Y(3S))}{\sigma(\text{direct } Y(1S))} = \frac{|\psi^{3S}(0)|^2}{|\psi^{1S}(0)|^2} \sim 0.34 \quad \frac{\sigma(\text{direct } Y(2S))}{\sigma(\text{direct } Y(1S))} = \frac{|\psi^{2S}(0)|^2}{|\psi^{1S}(0)|^2} \sim 0.45$$

Cross section ratio I

- Despite th. uncertainties, CSM predictions are **parameter free** !
- At LO in v^2 , one *de facto* **predicts direct** cross-section **ratios**
- Simple ratios of **Schrödinger wave function** at the origin:

$$\frac{\sigma(\text{direct } Y(3S))}{\sigma(\text{direct } Y(1S))} = \frac{|\psi^{3S}(0)|^2}{|\psi^{1S}(0)|^2} \sim 0.34 \quad \frac{\sigma(\text{direct } Y(2S))}{\sigma(\text{direct } Y(1S))} = \frac{|\psi^{2S}(0)|^2}{|\psi^{1S}(0)|^2} \sim 0.45$$

- $\sigma(Y(1S)(|y| < 2)) Br_{\ell\ell} \simeq 7.4 \text{ nb} \xrightarrow{50\% \text{ direct}} \sigma(\text{direct } Y(1S)) \sim 150 \text{ nb}$

CMS, PRD 83, 112004 (2011)

Cross section ratio I

- Despite th. uncertainties, CSM predictions are **parameter free** !
- At LO in v^2 , one *de facto* **predicts direct** cross-section **ratios**
- Simple ratios of **Schrödinger wave function** at the origin:

$$\frac{\sigma(\text{direct } Y(3S))}{\sigma(\text{direct } Y(1S))} = \frac{|\psi^{3S}(0)|^2}{|\psi^{1S}(0)|^2} \sim 0.34 \quad \frac{\sigma(\text{direct } Y(2S))}{\sigma(\text{direct } Y(1S))} = \frac{|\psi^{2S}(0)|^2}{|\psi^{1S}(0)|^2} \sim 0.45$$

- $\sigma(Y(1S)(|y| < 2)) Br_{\ell\ell} \simeq 7.4 \text{ nb} \xrightarrow{50\% \text{ direct}} \sigma(\text{direct } Y(1S)) \sim 150 \text{ nb}$

CMS, PRD 83, 112004 (2011)

- **Extrapolated** 3S direct yield: $0.34 \times 150 \text{ nb} \sim 50 \text{ nb}$

Cross section ratio I

- Despite th. uncertainties, CSM predictions are **parameter free** !
- At LO in v^2 , one *de facto* **predicts direct** cross-section **ratios**
- Simple ratios of **Schrödinger wave function** at the origin:

$$\frac{\sigma(\text{direct } Y(3S))}{\sigma(\text{direct } Y(1S))} = \frac{|\psi^{3S}(0)|^2}{|\psi^{1S}(0)|^2} \sim 0.34 \quad \frac{\sigma(\text{direct } Y(2S))}{\sigma(\text{direct } Y(1S))} = \frac{|\psi^{2S}(0)|^2}{|\psi^{1S}(0)|^2} \sim 0.45$$

- $\sigma(Y(1S)(|y| < 2)) Br_{\ell\ell} \simeq 7.4 \text{ nb} \xrightarrow{50\% \text{ direct}} \sigma(\text{direct } Y(1S)) \sim 150 \text{ nb}$

CMS, PRD 83, 112004 (2011)

- **Extrapolated** 3S direct yield: $0.34 \times 150 \text{ nb} \sim 50 \text{ nb}$

- $\sigma(Y(3S)(|y| < 2)) Br_{\ell\ell} \simeq 1.0 \text{ nb} \xrightarrow{100\% \text{ direct}} \sigma(\text{direct } Y(3S)) \sim 45 \text{ nb}$

CMS, PRD 83, 112004 (2011)

Cross section ratio I

- Despite th. uncertainties, CSM predictions are **parameter free** !
- At LO in v^2 , one *de facto* **predicts direct** cross-section **ratios**
- Simple ratios of **Schrödinger wave function** at the origin:

$$\frac{\sigma(\text{direct } \Upsilon(3S))}{\sigma(\text{direct } \Upsilon(1S))} = \frac{|\psi^{3S}(0)|^2}{|\psi^{1S}(0)|^2} \sim 0.34 \quad \frac{\sigma(\text{direct } \Upsilon(2S))}{\sigma(\text{direct } \Upsilon(1S))} = \frac{|\psi^{2S}(0)|^2}{|\psi^{1S}(0)|^2} \sim 0.45$$

- $\sigma(\Upsilon(1S)(|y| < 2)) Br_{\ell\ell} \simeq 7.4 \text{ nb} \xrightarrow{50\% \text{ direct}} \sigma(\text{direct } \Upsilon(1S)) \sim 150 \text{ nb}$

CMS, PRD 83, 112004 (2011)

- **Extrapolated** 3S direct yield: $0.34 \times 150 \text{ nb} \sim 50 \text{ nb}$



- $\sigma(\Upsilon(3S)(|y| < 2)) Br_{\ell\ell} \simeq 1.0 \text{ nb} \xrightarrow{100\% \text{ direct}} \sigma(\text{direct } \Upsilon(3S)) \sim 45 \text{ nb}$

CMS, PRD 83, 112004 (2011)



- **NEW**: the 3S yield likely not 100% direct

cf. $\chi_b(3P)$ observation by ATLAS PRL, 108, 152001 (2012)

Cross section ratio II





Cross section ratio II





- P_T dependence of cross section ratios:

Cross section ratio II





- P_T dependence of cross section ratios:
- Mass effects at low P_T : not incoded in the v^2 results: $M_{\text{NRQCD}}^{Y(nS)} = 2m_b$

Cross section ratio II



- P_T dependence of cross section ratios:
- Mass effects at low P_T : not incoded in the v^2 results: $M_{\text{NRQCD}}^{Y(nS)} = 2m_b$
- Feed-down: simple kinematical effect: $P_T^{\text{daughter}} \sim \frac{M^{\text{daughter}}}{M^{\text{mother}}} P_T^{\text{mother}}$



Cross section ratio II



- P_T dependence of cross section ratios:


- Mass effects at low P_T : not incoded in the v^2 results: $M_{\text{NRQCD}}^{Y(nS)} = 2m_b$
- Feed-down: simple kinematical effect: $P_T^{\text{daughter}} \sim \frac{M^{\text{daughter}}}{M^{\text{mother}}} P_T^{\text{mother}}$
- Harmless if $\frac{d\sigma}{dP_T} \propto P_T^{-n}$ with n fixed,

Cross section ratio II



- P_T dependence of cross section ratios:
 - Mass effects at low P_T : not incoded in the v^2 results: $M_{\text{NRQCD}}^{Y(nS)} = 2m_b$
 - Feed-down: simple kinematical effect: $P_T^{\text{daughter}} \sim \frac{M_{\text{daughter}}}{M_{\text{mother}}} P_T^{\text{mother}}$
 - Harmless if $\frac{d\sigma}{dP_T} \propto P_T^{-n}$ with n fixed,
 - harmful if n changes, esp. true at low P_T where $\frac{d\sigma}{dP_T}$ can be flat

Impact of P -waves




ChiGen: L. A. Harland-Lang and W. J. Stirling, <http://projects.hepforge.org/superchic/chigen.html>

NLO NRQCD: Y.-Q. Ma, K. Wang, K.-T. Chao, Phys. Rev. D83 (2011) 111503 (R)

The most important and overlooked theory paper on quarkonium physics in 2010 !

LHCb, Phys.Lett. B718 (2012) 431-440

Impact of P -waves



ChiGen: L. A. Harland-Lang and W. J. Stirling, <http://projects.hepforge.org/superchic/chigen.html>

NLO NRQCD: Y.-Q. Ma, K. Wang, K.-T. Chao, Phys. Rev. D83 (2011) 111503 (R)


The most important and overlooked theory paper on quarkonium physics in 2010 !

LHCb, Phys.Lett. B718 (2012) 431-440

- LHCb: first indication that the χ_c fraction increases

Note: NLO NRQCD does **not necessarily** mean “Colour Octet dominance”. At NLO, the Colour-Singlet and Colour-Octet transition yields depend **–for the P waves–** on the unphysical scale Λ_{NRQCD} and the NRQCD subtraction scheme

Impact of P -waves



ChiGen: L. A. Harland-Lang and W. J. Stirling, <http://projects.hepforge.org/superchic/chigen.html>

NLO NRQCD: Y.-Q. Ma, K. Wang, K.-T. Chao, Phys. Rev. D83 (2011) 111503 (R)

The most important and overlooked theory paper on quarkonium physics in 2010 !

LHCb, Phys.Lett. B718 (2012) 431-440


- LHCb: first indication that the χ_c fraction increases

Note: NLO NRQCD does **not necessarily** mean "Colour Octet dominance". At NLO, the Colour-Singlet and Colour-Octet transition yields depend **for the P waves** on the unphysical scale Λ_{NRQCD} and the NRQCD subtraction scheme

- About 40 % of $\Upsilon(1S)$ are from χ_b at the Tevatron

CDF, PRL 84, 2094 (2000).

Impact of P -waves



ChiGen: L. A. Harland-Lang and W. J. Stirling, <http://projects.hepforge.org/superchic/chigen.html>

NLO NRQCD: Y.-Q. Ma, K. Wang, K.-T. Chao, Phys. Rev. D83 (2011) 111503 (R)

The most important and overlooked theory paper on quarkonium physics in 2010 !


LHCb, Phys.Lett. B718 (2012) 431-440

- LHCb: first indication that the χ_c fraction increases

Note: NLO NRQCD does **not necessarily** mean “Colour Octet dominance”. At NLO, the Colour-Singlet and Colour-Octet transition yields depend **–for the P waves–** on the unphysical scale Λ_{NRQCD} and the NRQCD subtraction scheme

- About 40 % of $\Upsilon(1S)$ are from χ_b at the Tevatron
- At the LHC:

CDF, PRL 84, 2094 (2000).



A priori: no P_T dependence. However, the plot scales are different

LHCb JHEP 1211 (2012) 031


QCD corrections, feed-down and polarisation

B. Gong, J.X Wang, Phys. Rev. Lett. 100,232001,2008. P.Artoisenet, J.Campbell,JPL, F.Maltoni, F. Tramontano, Phys. Rev. Lett. 101,152001,2008
JPL, EPJC 61,693,2009. JPL, PLB 695,149,2011.

QCD corrections, feed-down and polarisation

B. Gong, J.X Wang, Phys. Rev. Lett. 100,232001,2008. P.Artoisenet, J.Campbell,JPL, F.Maltoni, F. Tramontano, Phys. Rev. Lett. 101,152001,2008
JPL, EPJC 61,693,2009. JPL, PLB 695,149,2011.


→ **Complete modification** of the CSM polarisation at NLO (also at NNLO*)



QCD corrections, feed-down and polarisation

B. Gong, J.X Wang, Phys. Rev. Lett. 100,232001,2008. P.Artoisenet, J.Campbell,JPL, F.Maltoni, F. Tramontano, Phys. Rev. Lett. 101,152001,2008
JPL, EPJC 61,693,2009. JPL, PLB 695,149,2011.

→ **Complete modification** of the CSM polarisation at NLO (also at NNLO*)




→ Polarisation from χ_Q Feed-down at NLO ?

QCD corrections, feed-down and polarisation

B. Gong, J.X Wang, Phys. Rev. Lett. 100,232001,2008. P.Artoisenet, J.Campbell,JPL, F.Maltoni, F. Tramontano, Phys. Rev. Lett. 101,152001,2008
JPL, EPJC 61,693,2009. JPL, PLB 695,149,2011.

→ **Complete modification** of the CSM polarisation at NLO (also at NNLO*)




→ Polarisation from χ_Q Feed-down at NLO ?

- If $\chi_Q \rightarrow {}^3S_1\gamma$ is E1: $\alpha_{from\chi_Q}^{max} = +1.00$ and $\alpha_{from\chi_Q}^{min} = -0.45$

QCD corrections, feed-down and polarisation

B. Gong, J.X Wang, Phys. Rev. Lett. 100,232001,2008. P.Artoisenet, J.Campbell,JPL, F.Maltoni, F. Tramontano, Phys. Rev. Lett. 101,152001,2008
JPL, EPJC 61,693,2009. JPL, PLB 695,149,2011.

→ **Complete modification** of the CSM polarisation at NLO (also at NNLO*)



→ Polarisation from χ_Q Feed-down at NLO ?


- If $\chi_Q \rightarrow {}^3S_1\gamma$ is E1: $\alpha_{from\chi_Q}^{max} = +1.00$ and $\alpha_{from\chi_Q}^{min} = -0.45$
- For the J/ψ :

JPL J. Phys. G 38 (2011) 124110

QCD corrections, feed-down and polarisation



B. Gong, J.X Wang, Phys. Rev. Lett. 100,232001,2008. P.Artoisenet, J.Campbell,JPL, F.Maltoni, F. Tramontano, Phys. Rev. Lett. 101,152001,2008
JPL EPJC 61,693,2009. JPL, PLB 695,149,2011.

→ **Complete modification** of the CSM polarisation at NLO (also at NNLO*)



→ Polarisation from χ_Q Feed-down at NLO ?

- If $\chi_Q \rightarrow {}^3S_1\gamma$ is E1: $\alpha_{from\chi_Q}^{max} = +1.00$ and $\alpha_{from\chi_Q}^{min} = -0.45$
- For the J/ψ :




JPL J. Phys. G 38 (2011) 124110

QCD corrections, feed-down and polarisation

B. Gong, J.X Wang, Phys. Rev. Lett. 100,232001,2008. P.Artoisenet, J.Campbell,JPL, F.Maltoni, F. Tramontano, Phys. Rev. Lett. 101,152001,2008
JPL, EPJ C 61,693,2009. JPL, PLB 695,149,2011.



→ **Complete modification** of the CSM polarisation at NLO (also at NNLO*)





→ Polarisation from χ_Q Feed-down at NLO ?

• If $\chi_Q \rightarrow {}^3S_1\gamma$ is E1: $\alpha_{from\chi_Q}^{max} = +1.00$ and $\alpha_{from\chi_Q}^{min} = -0.45$

• For the J/ψ :



• For the $Y(1S)$:



Part II

Associated Υ production

Associated production: J/ψ vs. Υ

- A number of associated-production channels proposed for J/ψ
 - $J/\psi + J/\psi$
 - $J/\psi + \gamma$
 - $J/\psi + c$ or $J/\psi + D$ or $J/\psi + lepton$
 - $J/\psi + Z$
 - $J/\psi + W$

Associated production: J/ψ vs. Υ

- A number of associated-production channels proposed for J/ψ
 - $J/\psi + J/\psi$
 - $J/\psi + \gamma$
 - $J/\psi + c$ or $J/\psi + D$ or $J/\psi + lepton$
 - $J/\psi + Z$
 - $J/\psi + W$
- Less studies for Υ
 - rates are usually lower
 - $J/\psi + c$ and $J/\psi + J/\psi$, sometimes motivated by intrinsic charm
Intrinsic bottom expected to be 10 times smaller

$\Upsilon + \text{prompt } \gamma$

- At high energy, 2 gluons in the initial states: no quark

$\Upsilon + \text{prompt } \gamma$

- At high energy, 2 gluons in the initial states: no quark
- The photon needs to be emitted by the b -quark loop

$\Upsilon + \text{prompt } \gamma$


- At high energy, 2 gluons in the initial states: no quark
- The photon needs to be emitted by the b -quark loop
- Gluon fragmentation associated with $C = +1$ octet ($^1S_0^{[8]}$ and $^3P_J^{[8]}$)

$\Upsilon + \text{prompt } \gamma$

- At high energy, 2 gluons in the initial states: no quark
- The photon needs to be emitted by the b -quark loop
- Gluon fragmentation associated with $C = +1$ octet ($^1S_0^{[8]}$ and $^3P_J^{[8]}$)
- CO rates may be **clearly lower if** $^1S_0^{[8]}$ and $^3P_J^{[8]}$ are indeed suppressed

Υ + prompt γ


- At high energy, 2 gluons in the initial states: no quark
- The photon needs to be emitted by the b -quark loop
- Gluon fragmentation associated with $C = +1$ octet ($^1S_0^{[8]}$ and $^3P_J^{[8]}$)
- CO rates may be **clearly lower** if $^1S_0^{[8]}$ and $^3P_J^{[8]}$ are indeed suppressed



JPL, PLB 679,340,2009.

Υ + prompt γ

- At high energy, 2 gluons in the initial states: no quark
- The photon needs to be emitted by the b -quark loop
- Gluon fragmentation associated with $C = +1$ octet ($^1S_0^{[8]}$ and $^3P_J^{[8]}$)
- CO rates may be **clearly lower** if $^1S_0^{[8]}$ and $^3P_J^{[8]}$ are indeed suppressed




JPL, PLB 679,340,2009.

- Clearly, **new info on CS vs CO w.r.t inclusive case !**

$\Upsilon + \text{prompt } \gamma$

- At high energy, 2 gluons in the initial states: no quark
- The photon needs to be emitted by the b -quark loop
- Gluon fragmentation associated with $C = +1$ octet ($^1S_0^{[8]}$ and $^3P_J^{[8]}$)
- CO rates may be **clearly lower** if $^1S_0^{[8]}$ and $^3P_J^{[8]}$ are indeed suppressed




JPL, PLB 679,340,2009.

- Clearly, **new info on CS vs CO w.r.t inclusive case !**
- Possible: see $(c, b) - jet + \gamma$ studies by D0 up to $P_T^\gamma \simeq 150 \text{ GeV} !$


D0, PRL102 (2009) 192002.

$\Upsilon + b$ -tagged jet




- $\Upsilon + b$: ~ 0.1 pb/GeV at the Tevatron, ~ 1 pb/GeV at the LHC (14 TeV)
- hard P_T spectrum w.r.t. the inclusive LO CSM


$\Upsilon + b$ -tagged jet




- $\Upsilon + b$: ~ 0.1 pb/GeV at the Tevatron, ~ 1 pb/GeV at the LHC (14 TeV)
- hard P_T spectrum w.r.t. the inclusive LO CSM
- $\Upsilon + b$: CSM vs. COM channels



$\Upsilon + b$ -tagged jet



- $\Upsilon + b$: ~ 0.1 pb/GeV at the Tevatron, ~ 1 pb/GeV at the LHC (14 TeV)
- hard P_T spectrum w.r.t. the inclusive LO CSM
- $\Upsilon + b$: CSM vs. COM channels
- Different topologies:
 - CSM: 1 b away, 1 b near(er)
 - COM: 2 b 's away (from a recoiling gluon)



$Q + \text{vector boson}$

- $\Upsilon + \text{vector boson}$

$Q + \text{vector boson}$

- $\Upsilon + \text{vector boson}$
 - 95% C.L. upper limits obtained with $\mathcal{L} = 83\text{pb}^{-1}$ by CDF

$Q + \text{vector boson}$

- $Y + \text{vector boson}$

- 95% C.L. upper limits obtained with $\mathcal{L} = 83\text{pb}^{-1}$ by CDF

$$\begin{aligned}\sigma[p\bar{p} \rightarrow Y(1S) + W^\pm] \times Br(Y(1S) \rightarrow \mu\mu) &< 2.3 \text{ pb} \\ \sigma[p\bar{p} \rightarrow Y(1S) + Z^0] \times Br(Y(1S) \rightarrow \mu\mu) &< 2.5 \text{ pb}\end{aligned}\tag{1}$$

CDF Collaboration, PRL. 90 (2003) 221803

- NRQCD predictions (Signal dominated by CO into χ_b)

$$\begin{aligned}\sigma[p\bar{p} \rightarrow Y(1S) + W^\pm] \times Br(Y(1S) \rightarrow \mu\mu) &\simeq 0.025 \text{ pb} \\ \sigma[p\bar{p} \rightarrow Y(1S) + Z^0] \times Br(Y(1S) \rightarrow \mu\mu) &\simeq 0.0075 \text{ pb}\end{aligned}\tag{2}$$

E. Braaten, J. Lee, and S. Fleming, PRD 60, 91501 (1999)

$Q + \text{vector boson}$

- $Y + \text{vector boson}$

- 95% C.L. upper limits obtained with $\mathcal{L} = 83\text{pb}^{-1}$ by CDF

$$\begin{aligned}\sigma[p\bar{p} \rightarrow Y(1S) + W^\pm] \times Br(Y(1S) \rightarrow \mu\mu) &< 2.3 \text{ pb} \\ \sigma[p\bar{p} \rightarrow Y(1S) + Z^0] \times Br(Y(1S) \rightarrow \mu\mu) &< 2.5 \text{ pb}\end{aligned}\tag{1}$$

CDF Collaboration, PRL. 90 (2003) 221803

- NRQCD predictions (Signal dominated by CO into χ_b)

$$\begin{aligned}\sigma[p\bar{p} \rightarrow Y(1S) + W^\pm] \times Br(Y(1S) \rightarrow \mu\mu) &\simeq 0.025 \text{ pb} \\ \sigma[p\bar{p} \rightarrow Y(1S) + Z^0] \times Br(Y(1S) \rightarrow \mu\mu) &\simeq 0.0075 \text{ pb}\end{aligned}\tag{2}$$

E. Braaten, J. Lee, and S. Fleming, PRD 60, 91501 (1999)

- CSM yield expected to be 300 times smaller (to be re-checked ...)

$Q + \text{vector boson}$

- $Y + \text{vector boson}$

- 95% C.L. upper limits obtained with $\mathcal{L} = 83\text{pb}^{-1}$ by CDF

$$\begin{aligned}\sigma[p\bar{p} \rightarrow Y(1S) + W^\pm] \times Br(Y(1S) \rightarrow \mu\mu) &< 2.3 \text{ pb} \\ \sigma[p\bar{p} \rightarrow Y(1S) + Z^0] \times Br(Y(1S) \rightarrow \mu\mu) &< 2.5 \text{ pb}\end{aligned}\tag{1}$$

CDF Collaboration, PRL. 90 (2003) 221803

- NRQCD predictions (Signal dominated by CO into χ_b)

$$\begin{aligned}\sigma[p\bar{p} \rightarrow Y(1S) + W^\pm] \times Br(Y(1S) \rightarrow \mu\mu) &\simeq 0.025 \text{ pb} \\ \sigma[p\bar{p} \rightarrow Y(1S) + Z^0] \times Br(Y(1S) \rightarrow \mu\mu) &\simeq 0.0075 \text{ pb}\end{aligned}\tag{2}$$

E. Braaten, J. Lee, and S. Fleming, PRD 60, 91501 (1999)

- CSM yield expected to be 300 times smaller (to be re-checked ...)
- With 1 fb^{-1} at $\sqrt{s} = 7 \text{ TeV}$ and a larger $(\mathcal{E} \times \mathcal{A})(Y)$, one should see events if CO's are at work

Q + vector boson

- Y + vector boson

- 95% C.L. upper limits obtained with $\mathcal{L} = 83\text{pb}^{-1}$ by CDF
$$\sigma[p\bar{p} \rightarrow Y(1S) + W^\pm] \times Br(Y(1S) \rightarrow \mu\mu) < 2.3 \text{ pb}$$
$$\sigma[p\bar{p} \rightarrow Y(1S) + Z^0] \times Br(Y(1S) \rightarrow \mu\mu) < 2.5 \text{ pb}$$

CDF Collaboration, PRL. 90 (2003) 221803

- NRQCD predictions (Signal dominated by CO into χ_b)

$$\sigma[p\bar{p} \rightarrow Y(1S) + W^\pm] \times Br(Y(1S) \rightarrow \mu\mu) \simeq 0.025 \text{ pb}$$
$$\sigma[p\bar{p} \rightarrow Y(1S) + Z^0] \times Br(Y(1S) \rightarrow \mu\mu) \simeq 0.0075 \text{ pb}$$

E. Braaten, J. Lee, and S. Fleming, PRD 60, 91501 (1999)

- CSM yield expected to be 300 times smaller (to be re-checked ...)
- With 1 fb^{-1} at $\sqrt{s} = 7 \text{ TeV}$ and a larger $(\mathcal{E} \times \mathcal{A})(Y)$, one should see events if CO's are at work

- $J/\psi + Z$ and $J/\psi + W$ recently computed at NLO in α_s

L.Gang *et al.* PRD83,014001,2011; JHEP02(2011)071

- $J/\psi|Y + Z$ at NLO in α_s + Polarisation

B.Gong *et al.* arXiv:1210.2430 [hep-ph] to appear in JHEP

$\Upsilon + Z$ cross sections



B. Gong, J.P. Lansberg, C. Lorcé, J.X. Wang, arXiv:1210.2430 [hep-ph] to appear in JHEP

- Rates similar for $\Upsilon + Z$ and $J/\psi + Z$!

$\Upsilon + Z$ cross sections

B. Gong, J.P. Lansberg, C. Lorcé, J.X. Wang, arXiv:1210.2430 [hep-ph] to appear in JHEP


- Rates similar for $\Upsilon + Z$ and $J/\psi + Z$!



$\Upsilon + Z$ cross sections

B. Gong, J.P. Lansberg, C. Lorcé, J.X. Wang, arXiv:1210.2430 [hep-ph] to appear in JHEP


- Rates similar for $\Upsilon + Z$ and $J/\psi + Z$!



- Mass effects ($m_c \leftrightarrow m_b$ less relevant because of m_Z)
- $|R(0)|^2$ is 10 times larger for Υ than for J/ψ
- Branching “only” 2.5 times smaller


$Y + Z$: Y polarisation

B. Gong, J.P. Lansberg, C. Lorcé, J.X. Wang, arXiv:1210.2430 [hep-ph] to appear in JHEP



$Y + Z$: Y polarisation


B. Gong, J.P. Lansberg, C. Lorcé, J.X. Wang, arXiv:1210.2430 [hep-ph] to appear in JHEP



- Y polarisation at LO and NLO are **similar**

$Y + Z$: Y polarisation


B. Gong, J.P. Lansberg, C. Lorcé, J.X. Wang, arXiv:1210.2430 [hep-ph] to appear in JHEP



- Y polarisation at LO and NLO are **similar**
- unlike the inclusive case
- not clear why: need for further investigation

$Y + Z$: Y polarisation

B. Gong, J.P. Lansberg, C. Lorcé, J.X. Wang, arXiv:1210.2430 [hep-ph] to appear in JHEP



- Y polarisation at LO and NLO are **similar**
- unlike the inclusive case
- not clear why: need for further investigation
- **CSM predictions** seem **robust** both for the yield and the polarisation

Part III

Υ in $p(d)A$ at RHIC and the LHC

Absorption: Υ vs. J/ψ

E.G. Ferreiro *et al.* arXiv:1110.5047 v4 [hep-ph]

σ_{abs}^{Υ} should be small \rightarrow nuclear PDF should play the major role

At RHIC:

Absorption: Υ vs. J/ψ

E.G. Ferreira *et al.* arXiv:1110.5047 v4 [hep-ph]

σ_{abs}^{Υ} should be small \rightarrow nuclear PDF should play the major role

At RHIC:

- at $y > 0$, $t_f = \gamma \times 0.4\text{fm} \gg r_{\text{Au}}$: pre-resonnant state exiting the nucleus

$$\sigma_{abs}^{b\bar{b}} \simeq 0.1 \sigma_{abs}^{c\bar{c}}$$

Absorption: Υ vs. J/ψ

E.G. Ferreira *et al.* arXiv:1110.5047 v4 [hep-ph]

σ_{abs}^{Υ} should be small \rightarrow nuclear PDF should play the major role

At RHIC:

- at $y > 0$, $t_f = \gamma \times 0.4\text{fm} \gg r_{Au}$: pre-resonnant state exiting the nucleus

$$\sigma_{abs}^{b\bar{b}} \simeq 0.1 \sigma_{abs}^{c\bar{c}}$$

- at $y < 0$, $t_f \leq r_{Au}$, fully formed in the nucleus: $\sigma_{abs}^{\Upsilon(2S)} \geq 4\sigma_{abs}^{\Upsilon(1S)}$

- Yet, equal suppression found by E772 in the backward region:

$$\sigma_{abs}^{\Upsilon(2S)} - \sigma_{abs}^{\Upsilon(1S)} \text{ small} \Rightarrow \sigma_{abs}^{\Upsilon(1S)} \text{ small}$$

Absorption: Υ vs. J/ψ

E.G. Ferreira *et al.* arXiv:1110.5047 v4 [hep-ph]

σ_{abs}^Y should be small \rightarrow nuclear PDF should play the major role

At RHIC:

- at $y > 0$, $t_f = \gamma \times 0.4\text{fm} \gg r_{Au}$: pre-resonnant state exiting the nucleus

$$\sigma_{abs}^{b\bar{b}} \simeq 0.1 \sigma_{abs}^{c\bar{c}}$$

- at $y < 0$, $t_f \leq r_{Au}$, fully formed in the nucleus: $\sigma_{abs}^{Y(2S)} \geq 4\sigma_{abs}^{Y(1S)}$


- Yet, equal suppression found by E772 in the backward region:

$$\sigma_{abs}^{Y(2S)} - \sigma_{abs}^{Y(1S)} \text{ small} \Rightarrow \sigma_{abs}^{Y(1S)} \text{ small}$$

At the LHC, the $b\bar{b}$ pair propagating in the nuclear matter (the Pb nucleus) is nearly always in a pre-resonnant (small) state


nPDF uncertainty on Υ production in $d\text{Au}$ collisions at RHIC

E.G. Ferreira *et al.* arXiv:1110.5047 v4 [hep-ph]



nPDF uncertainty on Υ production in $d\text{Au}$ collisions at RHIC


E.G. Ferreira *et al.* arXiv:1110.5047 v4 [hep-ph]



- Similar trend for the three nPDFs used (EKS98, EPS08, nDSg)

nPDF uncertainty on Υ production in d Au collisions at RHIC


E.G. Ferreira *et al.* arXiv:1110.5047 v4 [hep-ph]



- Similar trend for the three nPDFs used (EKS98, EPS08, nDSg)
- Uncertainty from the nPDFs: larger than our conservative estimate for that from the absorption x-section (0-1 mb)

nPDF uncertainty on Υ production in d Au collisions at RHIC


E.G. Ferreiro *et al.* arXiv:1110.5047 v4 [hep-ph]



- Similar trend for the three nPDFs used (EKS98, EPS08, nDSg)
- Uncertainty from the nPDFs: larger than our conservative estimate for that from the absorption x-section (0-1 mb)
- A priori, uncertainties smaller than for J/ψ
- But,

nPDF uncertainty on Υ production in $d\text{Au}$ collisions at RHIC

E.G. Ferreira *et al.* arXiv:1110.5047 v4 [hep-ph]



- Similar trend for the three nPDFs used (EKS98, EPS08, nDSg)
- Uncertainty from the nPDFs: larger than our conservative estimate for that from the absorption x-section (0-1 mb)
- A priori, uncertainties smaller than for J/ψ
- But, uncertainties in the EMC region are in fact large ...

Idea of a measurement for LHCb

E.G. Ferreira *et al.* arXiv:1110.5047 v4 [hep-ph]

- Cross section ratio for opposite rapidities in cms frame
[shift y_{lab} by -0.47]

Idea of a measurement for LHCb


E.G. Ferreira *et al.* arXiv:1110.5047 v4 [hep-ph]

- Cross section ratio for opposite rapidities in cms frame
[shift y_{lab} by -0.47]
- The physical interpretation (the meaning of a departure from unity)
does NOT depend on the pp yield

Idea of a measurement for LHCb

E.G. Ferreira *et al.* arXiv:1110.5047 v4 [hep-ph]

- Cross section ratio for opposite rapidities in cms frame
[shift y_{lab} by -0.47]
- The physical interpretation (the meaning of a departure from unity)
does NOT depend on the pp yield



Conclusions and Outlooks

- **LO** pQCD (CSM) reproduces the P_T -integrated yield:
relevant for heavy-ion studies: LO CSM is $gg \rightarrow Qg$

Conclusions and Outlooks

- **LO** pQCD (CSM) reproduces the P_T -integrated yield:
relevant for heavy-ion studies: LO CSM is $gg \rightarrow Qg$
- **LO** CSM fails as far as $d\sigma/dP_T$ is concerned

Conclusions and Outlooks

- **LO** pQCD (CSM) reproduces the P_T -integrated yield:
relevant for heavy-ion studies: LO CSM is $gg \rightarrow Qg$
- **LO** CSM fails as far as $d\sigma/dP_T$ is concerned
- Higher-QCD corrections open leading P_T channel: they are needed !

$2 \rightarrow 3, 2 \rightarrow 4$ channels

Conclusions and Outlooks

- **LO** pQCD (CSM) reproduces the P_T -integrated yield:
relevant for heavy-ion studies: LO CSM is $gg \rightarrow Qg$
- **LO** CSM fails as far as $d\sigma/dP_T$ is concerned
- Higher-QCD corrections open leading P_T channel: they are needed !
 $2 \rightarrow 3, 2 \rightarrow 4$ channels
- **Drawback**: large theoretical uncertainties...
Dominant contributions are known only at Born order ($gg \rightarrow Yggg$)

Conclusions and Outlooks

- **LO** pQCD (CSM) reproduces the P_T -integrated yield:
relevant for heavy-ion studies: LO CSM is $gg \rightarrow Qg$
- **LO** CSM fails as far as $d\sigma/dP_T$ is concerned
- Higher-QCD corrections open leading P_T channel: they are needed !
 $2 \rightarrow 3, 2 \rightarrow 4$ channels
- **Drawback**: large theoretical uncertainties...
Dominant contributions are known only at Born order ($gg \rightarrow Yggg$)
- In pA at RHIC, small nuclear effect on Y

Conclusions and Outlooks

- **LO** pQCD (CSM) reproduces the P_T -integrated yield:
relevant for heavy-ion studies: LO CSM is $gg \rightarrow Qg$
- **LO** CSM fails as far as $d\sigma/dP_T$ is concerned
- Higher-QCD corrections open **leading** P_T channel: they are needed !
 $2 \rightarrow 3, 2 \rightarrow 4$ channels
- **Drawback**: large theoretical uncertainties...
Dominant contributions are known only at Born order ($gg \rightarrow Yggg$)
- In pA at RHIC, small nuclear effect on Y
- But **backward- Y** sensitive to gluon EMC effect, which can be large
[hint from the data]

Conclusions and Outlooks

- **LO** pQCD (CSM) reproduces the P_T -integrated yield:
relevant for heavy-ion studies: LO CSM is $gg \rightarrow Qg$
- **LO** CSM fails as far as $d\sigma/dP_T$ is concerned
- Higher-QCD corrections open **leading** P_T channel: they are needed !

$2 \rightarrow 3, 2 \rightarrow 4$ channels

- **Drawback**: large theoretical uncertainties. . .
Dominant contributions are known only at Born order ($gg \rightarrow Yggg$)
- In pA at RHIC, small nuclear effect on Y
- But **backward- Y** sensitive to gluon EMC effect, which can be large
[hint from the data]
- Suggestion for the pPb data: forward-backward ratio
[pp reference irrelevant]